

Decreased visual acuity secondary to celiac disease

Baixa acuidade visual secundária à doença celíaca

Thiago Gonçalves dos Santos Martins¹, Ana Luiza Fontes de Azevedo Costa²



Cranial computed tomography showing posterior cortical-subcortical calcifications in the occipital and occipitoparietal regions

The male patient was a 24-year-old student who was a native of São Paulo (SP). The onset of left hemibody numbness associated with decreased strength and difficulty speaking was 15 days ago. It progressed with decreased bilateral visual acuity. After 90 minutes, the patient presented with photophobia and severe holocranial headaches, which were unresponsive to common analgesics. The headaches improved after analgesic medication, but the visual change persisted. For his past medical history, the patient has had recurrent episodes of “sparkles” in his vision that

progressed to headaches since he was 12 years of age. He was evaluated by a neurologist at that time and showed changes in the central nervous system, but no specific diagnosis was indicated. The patient reported the occurrence of four additional episodes over the past year that were similar to the current episode, although the intensity of the headaches was lower. In his family history, there have been cases of migraine headaches. Since the last episode, the patient has been taking valproic acid and dexamethasone. An ophthalmologic examination showed a best-corrected

¹ Department of Ophthalmology, Universidade Federal de São Paulo – UNIFESP, São Paulo (SP), Brazil.

² Hospital Federal dos Servidores do Estado do Rio de Janeiro – HFSE, Rio de Janeiro (RJ), Brazil.

Corresponding author: Thiago Gonçalves dos Santos Martins – Universidade Federal de São Paulo – UNIFESP – Departamento de Oftalmologia – Rua Botucatu, 821 – Zip code: 04023-062 – São Paulo (SP), Brazil – Phone: (55 11) 96456-2182 – Email: thiagogsmartins@yahoo.com.br

Received on: March 2, 2012 – Accepted on: Aug 23, 2012

visual acuity of 0.1 in both eyes. The remainder of the examination consisted of external ocular motility, biomicroscopy, Goldmann appplanation tonometry and neuromotor reflexes, which were all unchanged in both eyes. A cranial computerized tomography scan was performed, which showed posterior cortical-subcortical calcifications in the occipital and occipitoparietal regions. The patient was then referred for an evaluation to the gastroenterology service in the hospital, where he underwent a gastrointestinal endoscopy including a small bowel biopsy and received a definitive diagnosis of celiac disease. After eight months on a gluten-free diet, the patient showed improved visual acuity (1.0 in both eyes).

In 2004, Zelnik et al.⁽¹⁾ identified the co-existence of neurological symptoms in 51.4% of patients in a pediatric population with celiac disease compared to 19.9% in the control group. In a case report of a 4-year-old patient presenting with epileptic seizures, Lea et al.⁽²⁾ documented the progressive appearance of bilateral parietal intracranial calcifications and a subsequent diagnosis of celiac disease, which resulted in adequate seizure control without the need for medication after instituting dietary changes. The description of the triad of bilateral parietal calcifications, epilepsy and celiac disease is already widely reported in the literature⁽³⁻¹⁰⁾. Pfaender et al.⁽⁴⁾ reported the cases of three patients whose initial symptoms of disease were the occurrence of various visual symptoms, which were sometimes followed by significant headaches/migraines and the identification of bilateral parietal intracranial calcifications, with a subsequent diagnosis of celiac disease. This description reinforces the initial proposal that the occurrence of paroxysmal visual manifestations is the primary characteristic of epilepsy arising from the occipital lobe, despite being a rare focal form. The

three patients showed an improvement of symptoms after instituting a restrictive diet and antiepileptic medication.

The mechanisms involved in the onset of intracranial calcifications specifically localized to the bilateral parietal region have not yet been elucidated. *Post-mortem* analysis of brain tissue from some patients drives the suspicions that immune-mediated phenomena may alter the cerebral vascular permeability, favoring the localized deposition of calcium^(3,6).

REFERENCES

1. Zelnik N, Pacht A, Obeid R, Lerner A. Range of neurologic disorders in patients with celiac disease. *Pediatrics*. 2004;113(6):1672-6.
2. Lea ME, Harbord M, Sage MR. Bilateral occipital calcification associated with celiac disease, folate deficiency, and epilepsy. *AJNR*. 1995;16(7):1498-500.
3. Magaudda A, Dalla Bernardina B, De Marco P, Sfaello Z, Longo M, Colamaria V, et al. Bilateral occipital calcification, epilepsy and coeliac disease: clinical and neuroimaging features of a new syndrome. *J Neurol Neurosurg Psychiatry*. 1993;56(8):885-9.
4. Pfaender M, D'Souza W J, Trost N, Litewka L, Paine M, Cook M. Visual disturbances representing occipital lobe epilepsy in patients with cerebral calcifications and coeliac disease: a case series. *J Neurol Neurosurg Psychiatry*. 2004;75(11):1623-5.
5. Murray JA. The widening spectrum of celiac disease. *Am J Clin Nutr*. 1999; 69(3):354-65.
6. Bernasconi A, Bernasconi N, Andermann F, Dubeau FG, Guberman A, Gobbi G, et al. Celiac disease, bilateral occipital calcifications and intractable epilepsy: Mechanisms of seizure origin. *Epilepsia*. 1998;39(3):300-6.
7. Souza Santos CH, Brandão de Almeida IL, Gomes MDC, Serafim A, Pereira MM, Muszkat M, et al. Calcificação intracraniana occipital bilateral, epilepsia e doença celíaca: relato de caso. *Arq Neuropsiquiatr*. 2002;60(3-B):840-3.
8. Siqueira Neto JI, Vieira Costa ACL, Magalhães FG, Silva GS. Neurological manifestations of celiac disease. *Arq Neuropsiquiatr*. 2004;62(4):969-72.
9. Taylor I, Scheffer IE, Berkovic SF. Occipital epilepsies: identification of specific and newly recognized syndromes. *Brain*. 2003;126(Pt 4):753-69.
10. Palheta Neto FX, Vieira Junior MA, Ximenes LS, Jacob CCS, Rodrigues Junior AG, Palheta ACP. Aspectos clínicos da síndrome de Sturge-Weber. *Int Arch Otorhinolaryngol*. 2008;12(4):565-70.